



THE PRIMARY CARE COMPANION FOR CNS DISORDERS

Supplementary Material

Article Title: Posttraumatic Stress Disorder Symptoms, Functioning, and Suicidal Ideation in US Military Veterans: A Symptomics Approach

Authors: Lorig K. Kachadourian, PhD^{a,b,*}; Ilan Harpaz-Rotem, PhD^{a,b,c}; Jack Tsai, PhD^{a,b,c}; Steven M. Southwick, MD^{a,b}; and Robert H. Pietrzak, PhD, MPH^{a,b}

DOI Number: <https://doi.org/10.4088/PCC.18m02402>

List of Supplementary Material for the article

1. Supplementary Table 1. Results of Regression Analyses of Heterogeneity Models Evaluating Associations Between Individual PTSD Symptoms and Measures of Functioning, Quality of Life, and Suicidal Ideation

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Table 1. Results of Regression Analyses of Heterogeneity Models Evaluating Associations Between Individual PTSD Symptoms and Measures of Functioning, Quality of Life, and Suicidal Ideation

	Physical Functioning	Mental Functioning	Cognitive Functioning	Quality of Life	Suicidal Ideation
Mean (SD) or n (%)	46.7 (9.4)	52.6 (8.6)	87.7 (17.5)	52.4 (10.4)	120 (9.1%)
R^2	0.19	0.58	0.63	0.43	0.41
	β	β	β	β	OR
Unwanted memories	-0.01	-0.13***	0.03	-0.04	1.24
Nightmares	-0.09	-0.01	-0.13***	0.04	1.05
Flashbacks	0.06	0.07	0.01	0.01	0.66
Emotional reactivity to trauma	0.02	0.02	-0.01	0.03	0.91
Physical reactivity to trauma	-0.06	-0.10**	-0.15***	0.00	1.33
Avoiding thoughts and feelings	-0.08*	0.04	-0.02	-0.03	0.75
Avoiding reminders	0.02	0.02	0.05	0.01	0.91
Psychogenic amnesia	-0.11***	0.00	-0.10***	-0.05*	1.19
Negative thoughts	-0.03	-0.17***	-0.07*	-0.11**	1.55**
Exaggerated blame	0.00	-0.05	-0.01	-0.05	1.08
Negative affect	0.07	0.00	-0.07*	0.05	1.37
Loss of interest	-0.19***	-0.03	-0.01	-0.12**	0.86
Feeling isolated	0.06	-0.07	0.03	-0.09*	0.79
Difficulty experiencing positive affect	0.00	-0.13***	-0.11***	-0.16***	1.81**
Irritability/aggression	0.04	-0.11**	-0.07*	-0.03	1.39
Risky or destructive behavior	0.04	-0.01	0.01	0.01	1.46*
Hypervigilance	-0.06	0.07	0.02	0.04	0.77
Heightened startle reaction	-0.09*	-0.04	-0.10***	-0.03	1.05
Concentration difficulties	-0.04	-0.23***	-0.41***	-0.16***	1.34
Sleep difficulties	-0.14***	-0.04*	-0.05*	-0.14***	1.18
Age	-0.12***	0.12***	0.07**	0.07**	0.99
Male sex	0.04	0.05*	-0.06**	-0.01	1.04

Note. OR=odds ratio.

Significant association with dependent variable, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.